

Erectile  
Dysfunction

**Hospital discharges:** The estimated number of hospital discharges among men aged 20 years or older with a discharge diagnosis of "hyperplasia of the prostate":

(2001): 402,000 discharges <sup>9</sup>

(2000): 398,000 discharges <sup>10</sup>

**Doctor visits:** The estimated number of doctor visits by men aged 20 years or older with "hyperplasia of the prostate" as a listed diagnosis:

(2000): 6.4 million visits <sup>11</sup>

(1999): 6.0 million visits <sup>12</sup>

#### Erectile dysfunction (impotence)

**Prevalence** (2000-2002): 20 percent to 46 percent of men aged 40-69 years in community surveys self-reported moderate or complete erectile dysfunction. <sup>16-18</sup>

**Incidence** (2000): 26 new cases per 1,000 population of men 40-69 years of age <sup>19</sup>

#### Prostate cancer

**Incidence** (1996-2000): 975 cases per 100,000 population for men 65 years or older; 57 cases per 100,000 population for men under 65 years old <sup>20</sup>

**Mortality** (1996-2000): 245 deaths per 100,000 population for men 65 years or older; 2 deaths per 100,000 population for men under 65 years old <sup>20</sup>

**Lifetime prevalence** (2000): The number of living men who have had a diagnosis of prostate cancer is 1.6 million.

[Top]

#### Sources

1. K/DOQI Clinical Practice Guidelines for Chronic Kidney Disease: Evaluation, Classification, and Stratification. *American Journal of Kidney Disease*. 2002;39(2, Suppl. 1):S1-S266.
2. Summary of notifiable diseases, United States, 2001. *Morbidity and Mortality Weekly Report (MMWR)*. 2003;50